arXiv:1504.04573v1 [math.GT] 17 Apr 2015

REPRESENTATIONS OF THE KAUFFMAN SKEIN ALGEBRA OF SMALL SURFACES

NURDIN TAKENOV

Abstract. We prove a uniqueness result for finite-dimensional representations of the Kauffman skein algebra $\mathcal{S}_A(S)$ of a surface S, when A is a root of unity and when the surface S is a sphere with at most four punctures or a torus with at most one puncture. We show that, if two irreducible representations of $\mathcal{S}_A(S)$ have the same classical shadow and the same puncture invariants, and if this classical shadow is sufficiently generic in the character variety $\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$, then the two representations are isomorphic.

Let S be an oriented surface (without boundary) with finite topological type. The *Kauffman skein algebra* $\mathcal{S}_A(S)$ of S is a certain quantization of the character variety

$$\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S) = \{\text{group homomorphisms } r\colon \pi_1(S) \to \mathrm{SL}_2(\mathbb{C})\} /\!\!/ \mathrm{SL}_2(\mathbb{C})$$

depending on a parameter $A = \mathrm{e}^{\pi \mathrm{i} \hbar} \in \mathbb{C} - \{0\}$. The elements of $\mathcal{S}_A(S)$ are represented by linear combinations of framed links in the thickened surface $S \times [0,1]$, considered modulo certain relations; see [10, 6, 9] and §1.

In [2, 3, 4], Bonahon and Wong consider finite-dimensional representations of the skein algebra $\mathcal{S}_A(S)$ when A is a root of unit, construct invariants of irreducible representations, and show that any set of invariant is realized. The purpose of the current paper is to provide a uniqueness component to their results when the surface S is small.

The precise result of [2] is the following. The setup requires A^2 to be an N–root of unity with N odd, and we will restrict attention to the case where $A^N = -1$ as results are easier to state; because N is odd, this case is equivalent to requiring that A be an N–root of -1. The case $A^N = +1$ can be deduced from the case $A^N = -1$ by using spin structures on the surface; see [1, 2].

By definition of the geometric invariant theory quotient involved in the definition of $\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$, two homomorphisms $r, r'\colon \pi_1(S) \to \mathrm{SL}_2(\mathbb{C})$ define the same element of $\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$ if and only if they associate the same trace $\mathrm{Tr}\, r(\gamma) = \mathrm{Tr}\, r'(\gamma)$ to each element $\gamma \in \pi_1(S)$.

Theorem 1 ([2]). *Suppose that A is a primitive N-root of -1 with N odd, and let $\rho\colon \mathcal{S}_A(S) \to \mathrm{End}(V)$ be an irreducible finite-dimensional representation of the Kauffman skein algebra. Let $T_N(x)$ be the N-th normalized Chebyshev polynomial of the first kind, defined by the trigonometric equality that $\cos N\theta = \frac{1}{2} T_N(2\cos\theta)$.*

(1) *There exists a unique character $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$ such that*

$$T_N\big(\rho([K])\big) = -\big(\mathrm{Tr}\ r(K)\big)\,\mathrm{Id}_V$$

in $\mathrm{End}(V)$ for every framed knot $K \subset S \times [0,1]$ whose projection to S has no crossing and whose framing is vertical.

Date: May 9, 2015.

This research was partially supported by grants DMS-1105402 and DMS-1406559 from the U.S. National Science Foundation.

(2) *Let* P_k *be a small simple loop going around the* k*-th puncture of* S*, and consider it as a knot in* $S \times [0,1]$ *with vertical framing. Then there exists a number* $p_k \in \mathbb{C}$ *such that* $\rho\big([P_k]\big) = p_k \,\mathrm{Id}_V$*.*

(3) *The number* p_k *of* (2) *is related to the character* $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$ *of* (1) *by the property that* $T_N(p_k) = -\mathrm{Tr}\; r(P_k)$*.*

The character $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$ is the *classical shadow* of the representation ρ, while the numbers p_k are its *puncture invariants.* Bonahon and Wong also show in [3, 4] that every character $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$ and set of numbers $p_k \in \mathbb{C}$ associated to the punctures of S that satisfy the relation (3) of Theorem 1 can be realized as the set of invariants of an irreducible representation of $\mathcal{S}_A(S)$.

We address the following conjecture which asserts that, generically, an irreducible representation of $\mathcal{S}_A(S)$ should be determined by its classical shadow and its puncture invariants.

Conjecture 2 (Generic Uniqueness Conjecture)**.** *There exists a Zarisky dense open subset* $\mathcal{U} \subset \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$ *such that, if two irreducible representations* ρ, $\rho' \colon \mathcal{S}_A(S) \to \mathrm{End}(V)$ *have the same classical shadow and the same puncture invariants, and if the classical shadow* r *belongs to this dense subset* $\mathcal{U} \subset \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$*, then the representations* ρ *and* ρ' *are isomorphic.*

Our main result is the following.

Theorem 3. *The Generic Uniqueness Conjecture* 2 *holds when the surface* S *is a sphere with at most* 4 *punctures, or a torus with* 0 *or* 1 *puncture.*

The cases of the sphere with at most 3 punctures are essentially trivial; see §3. When S is the one-puncture torus, we rely in §4 on a presentation of $\mathcal{S}_A(S)$ given by Bullock-Przytycki [7] and on the systematic classification of all irreducible representations of the corresponding algebra by Havlíček-Pošta [8]; it is however non-trivial to relate the invariants of [8] to those of Theorem 1. The case of the unpunctured torus follows from the one-puncture case; see §5. For the four-puncture sphere, we again use in §6 a presentation of $\mathcal{S}_A(S)$ exhibited in [7], and extend to this context the arguments that we had developed for the one-puncture torus; however, the situation is very significantly more complicated than for the one-puncture torus.

The analysis of [8] provides many different representations of the skein algebra of the one-puncture torus whose classical shadow is the trivial character. As a consequence, Conjecture 2 cannot hold without the genericity hypothesis.

1. The Kauffman skein algebra

Let S be an oriented surface of finite topological type, without boundary. This means that $S = S_{g,p}$ is obtained by removing p points from a closed oriented surface $\bar{S}$ of genus g. We consider *framed links* in the thickened surface $S \times [0,1]$, namely unoriented 1-dimensional submanifolds $K \subset S \times [0,1]$ endowed with a continuous choice of a vector transverse to K at each point of K. A *framed knot* is a framed link that is connected.

The following definition provides a convenient way to describe a framing, in particular when representing a link by a picture. If the projection of $K \subset S \times [0,1]$ to S is an immersion, the *vertical framing* for K is the framing that is everywhere parallel to the $[0,1]$ factor and points towards 1.

The *framed link algebra* $\mathcal{K}(S)$ is the vector space over $\mathbb{C}$ freely generated by the isotopy classes of all framed links $K \subset S \times [0,1]$. This vector space $\mathcal{K}(S)$ can be endowed with a multiplication, where the product of K_1 and K_2 is defined as the framed link $K_1 \cdot K_2 \subset$

$S \times [0,1]$ that is the union of K_1 rescaled in $S \times [0, \frac{1}{2}]$ and K_2 rescaled in $S \times [\frac{1}{2}, 1]$. In other words, the product $K_1 \cdot K_2$ is defined by superposition of the framed links K_1 and K_2. This *superposition operation* is compatible with isotopies, and therefore provides a well-defined algebra structure on $\mathcal{K}(S)$.

Three framed links K_1, K_0 and K_∞ in $S \times [0,1]$ form a *Kauffman triple* if the only place where they differ is above a small disk in S, where they are as represented in Figure 1 (as seen from above) and where the framing is vertical and pointing upwards (namely the framing is parallel to the $[0,1]$ factor and points towards 1).



FIGURE 1. A Kauffman triple

For $A \in \mathbb{C} - \{0\}$, the *Kauffman skein algebra* $\mathcal{S}_A(S)$ is the quotient of the framed link algebra $\mathcal{K}(S)$ by the linear subspace generated by:

(1) all elements $K_1 - A^{-1}K_0 - AK_\infty \in \mathcal{K}(S)$ as (K_1, K_0, K_∞) ranges over all Kauffman triples;
(2) the element $\bigcirc + (A^2 + A^{-2})\varnothing$ where $\bigcirc \subset S \times [0,1]$ is an unknot projecting to asimple loop in S and endowed with the vertical framing, and where $\varnothing$ is the empty link.

The superposition operation descends to a multiplication in $\mathcal{S}_A(S)$, and endows $\mathcal{S}_A(S)$ with the structure of an algebra. The class $[\varnothing]$ of the empty link is an identity element in $\mathcal{S}_A(S)$, and we usually identify it to the scalar 1.

An element $[K] \in \mathcal{S}_A(S)$ represented by a framed link $K \subset S \times [0,1]$ is a *skein* in S.

Throughout the article, we will assume that the parameter A is a primitive N–root of -1 with N odd; for instance, $A = \mathrm{e}^{\pi \mathrm{i}/N}$. Because N is odd, this implies that A^2 and A^4 are primitive N–roots of unity, a property that is frequently used in the article.

2. CHEBYSHEV POLYNOMIALS

The *normalized n-th Chebyshev polynomial of the first kind* is the polynomial $T_n(x)$ such that $\operatorname{Tr} M^n = T_n(\operatorname{Tr} M)$ for every $M \in \mathrm{SL}_2(\mathbb{C})$. It can be inductively computed by the recurrence relation that $T_n(x) = xT_{n-1}(x) - T_{n-2}(x)$, combined with the initial conditions $T_0(x) = 2$ and $T_1(x) = x$.

The polynomial $T_n(x)$ contains only even powers of x when n is even, and only odd powers of x when n is odd. Also, applying the relation $\operatorname{Tr} M^n = T_n(\operatorname{Tr} M)$ to a rotation matrix gives the trigonometric identity that $2\cos n\theta = T_n(2\cos\theta)$.

We will use the following computations.

Lemma 4.

(a) $T_n(a + a^{-1}) = a^n + a^{-n}$;
(b) $T_N(x) - a^N - a^{-N} = \prod_{k=1}^{N} \left(x - aA^{2k} - a^{-1}A^{-2k}\right)$.

Proof. Property (a) is a simple application of the relation $\operatorname{Tr} M^n = T_n(\operatorname{Tr} M)$ to a matrix $M \in \mathrm{SL}_2(\mathbb{C})$ with eigenvalues a, a^{-1}.

For (b), note that (a) implies that the solutions of the equation $T_N(x) = a^N + a^{-N}$ are the numbers $x = b + b^{-1}$ with $b^N = a^N$. Since A^2 is a primitive N-root of unity, these are the numbers of the form $x = aA^{2k} + a^{-1}A^{-2k}$. The equality then follows from the fact that the highest degree term of $T_N(x)$ is x^N. □

3. THE SPHERE WITH AT MOST THREE PUNCTURES

This case is essentially trivial. Indeed, in the sphere with at most 3 punctures, every simple closed curve is isotopic, either to a trivial knot, or to a simple loop P_k going around one of the punctures. It follows that the algebra $\mathcal{S}_A(S)$ is generated by the skeins $[P_k]$, so that every irreducible representation $\rho\colon \mathcal{S}_A(S) \to \mathrm{End}(V)$ is completely determined by its puncture invariants p_k (which also determine the classical shadow).

4. THE ONE-PUNCTURE TORUS $S_{1,1}$

This section is devoted to the case of the one-puncture torus $S_{1,1}$.

4.1. **A presentation for the skein algebra $\mathcal{S}_A(S_{1,1})$.** We will use the presentation of $\mathcal{S}_A(S_{1,1})$ given by Bullock and Przytycki in [7].

Identify the punctured torus $S_{1,1}$ to the quotient space obtained from a square $[0,1] \times [0,1]$ by removing the four corners and gluing together opposite sides in the usual fashion. Let X_1 be the closed curve that is the image in $S_{1,1}$ of the vertical line segment $\{\frac{1}{2}\} \times [0,1]$, let X_2 be the image of the horizontal line segment $[0,1] \times \{\frac{1}{2}\}$, and let X_3 be the closed curve that is the image of the two slope 1 line segments respectively parametrized by $t \mapsto (t+\frac{1}{2}, t)$ and $t \mapsto (t, t+\frac{1}{2})$ with $0 \leq t \leq \frac{1}{2}$. See Figure 2.

Also, let $P \subset S_{1,1}$ be a simple loop going around the puncture.



FIGURE 2. Curves on the one-puncture torus

Proposition 5 ([7])**.** *The Kauffman skein algebra $\mathcal{S}_A(S_{1,1})$ admits a presentation by generators X_1, X_2 and X_3 and relations*

$$\begin{aligned} AX_1X_2 - A^{-1}X_2X_1 &= (A^2 - A^{-2})X_3 \\ AX_2X_3 - A^{-1}X_3X_2 &= (A^2 - A^{-2})X_1 \\ AX_3X_1 - A^{-1}X_1X_3 &= (A^2 - A^{-2})X_2 \end{aligned}$$

where the X_i are represented by the closed curves indicated above, endowed with vertical framing.

If the loop P going around the puncture is also endowed with the vertical framing, the corresponding element of $\mathcal{S}_A(S_{1,1})$ is equal to

$$P = AX_1X_2X_3 - A^2X_1^2 - A^{-2}X_2^2 - A^2X_3^2 + A^2 + A^{-2}.$$

This element is central in $\mathcal{S}_A(S_{1,1})$. □

The above presentation is also a presentation of another algebra, the algebra $\mathrm{U}_q'(\mathfrak{so}_3)$ whose irreducible representations were classified by Havlíček and Pošta [8]. We will heavily rely on the arguments of [8], while adapting them to our goals.

4.2. **Reconstructing an irreducible representation of $\mathcal{S}_A(S_{1,1})$.**

Theorem 6. *Let $\rho\colon \mathcal{S}_A(S_{1,1}) \to End(V)$ be an irreducible representation with classical shadow $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{1,1})$ and puncture invariant $p \in \mathbb{C}$. Suppose that*

$$\begin{aligned} &\mathrm{Tr}\, r(X_3) \neq \pm 2 \\ \text{and } &\mathrm{Tr}\, r(X_1)\mathrm{Tr}\, r(X_2)\mathrm{Tr}\, r(X_3) + \mathrm{Tr}\, r(X_1)^2 + \mathrm{Tr}\, r(X_2)^2 \neq 0 \end{aligned}$$

for the curves X_1, X_2, X_3 of §4.1*. Then, up to isomorphism, the representation $\rho\colon \mathcal{S}_A(S_{1,1}) \to End(V)$ is completely determined by r and p. In addition, V has dimension N.*

Theorem 6 proves our Generic Uniqueness Theorem 3 in the case of the one-puncture torus $S_{1,1}$, since its hypotheses describe a Zarisky dense open subset of the character variety $\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S)$. We will late provide a slightly more general result, Theorem 15.

The proof of Theorem 6 will take a while, and we will split it into several lemmas. We assume that the hypotheses of Theorem 6 hold, for the remainder of this section.

For notational convenience, set $t_i = -\mathrm{Tr}\, r(X_i)$ for $i = 1, 2, 3$, so that $T_N\big(\rho(X_i)\big) = t_i\,\mathrm{Id}_V$.The hypotheses of Theorem 6 are then that $t_3 \neq \pm 2$ and $t_1t_2t_3 + t_1^2 + t_2^2 \neq 0$.

The numbers t_1, t_2, t_3 and the puncture invariant p are related by the following equation.

Lemma 7.

$$T_N(p) = -t_1t_2t_3 - t_1^2 - t_2^2 - t_3^2 + 2$$

Proof. This is a consequence of the special case where $N = 1$ and $A = -1$. Indeed, an observation of Bullock [5] (see also [9]) shows that the character $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{1,1})$ uniquely determines a homomorphism $\Theta_r\colon \mathcal{S}_{-1}(S_{1,1}) \to \mathbb{C}$ by the property that $\Theta_r\big([K]\big) = -\mathrm{Tr}\, r(K)$ for every framed knot $K \subset S_{1,1} \times [0,1]$.

In the special case $A = -1$ considered, the second half of Proposition 5 states that the elements X_1, X_2, X_3, $P \in \mathcal{S}_{-1}(S_{1,1})$ satisfy the relation

$$P = -X_1X_2X_3 - X_1^2 - X_2^2 - X_3^2 + 2$$

in $\mathcal{S}_{-1}(S_{1,1})$. Applying the homomorphism Θ_r on both sides of the equation, and remembering that $\Theta_r(X_i) = -\operatorname{Tr} r(X_i) = t_i$ and $\Theta_r(P) = -\operatorname{Tr} r(P) = T_N(p)$, then provides the equality sought. $\square$

Choose a number $x_3 \in \mathbb{C}$ such that $t_3 = x_3^N + x_3^{-N}$.

Because $T_N\big(\rho(X_3)\big) = t_3 \operatorname{Id}_V$, all possible eigenvalues λ of $\rho(X_3)$ are such that $T_N(\lambda) = t_3 = x_3^N + x_3^{-N}$, and therefore equal to one of the numbers $\lambda_k = x_3 A^{2k} + x_3^{-1} A^{-2k}$ by Lemma 4(a). Since $t_3 \neq \pm 2$ by hypothesis, the numbers λ_k with $k = 1, 2, \dots, N$ are distinct. It is convenient to consider all integer indices k, so that $\lambda_{k+N} = \lambda_k$.

Define $V_k = \{v \in V; \rho(X_3)v = \lambda_k v\}$. Namely, V_k is the eigenspace corresponding to λ_k if λ_k is really an eigenvalue of $\rho(X_3)$, and is 0 otherwise.

Our key tool is provided by the operators $\mathcal{U}_k$ and $\mathcal{D}_k \in \operatorname{End}(V)$ defined by

$$\begin{aligned}\mathcal{U}_k &= A\rho(X_1) - x_3 A^{2k}\rho(X_2)\\ \mathcal{D}_k &= A\rho(X_1) - x_3^{-1} A^{-2k}\rho(X_2).\end{aligned}$$

These "up" and "down" operators $\mathcal{U}_k$ and $\mathcal{D}_k$ are borrowed from [8]. What makes them so useful is the following property.

Lemma 8. *The operator $\mathcal{U}_k$ sends the subspace V_k to V_{k+1}, and $\mathcal{D}_k$ sends V_k to V_{k-1}.*

Proof. We want to show that $\rho(X_3)\mathcal{U}_k v_k = \lambda_{k+1}\mathcal{U}_k v_k$ for every $v_k \in V_k$. For this, expand

$$\rho(X_3)\mathcal{U}_k v_k = \rho(X_3)\left(A\rho(X_1) - x_3 A^{2k}\rho(X_2)\right) v_k = \left(A\rho(X_3X_1) - x_3 A^{2k}\rho(X_3X_2)\right) v_k.$$

Using the relations of Proposition 5,

$$\begin{aligned}\rho(X_3)\mathcal{U}_k v_k &= (A\rho(X_3X_1) - x_3A^{2k}\rho(X_3X_2))v_k\\ &= A\rho(A^{-2}X_1X_3 + A^{-1}(A^2 - A^{-2})X_2)v_k\\ &\qquad - x_3A^{2k}\rho(A^2X_2X_3 - A(A^2 - A^{-2})X_1)v_k\\ &= A^{-1}\rho(X_1)\rho(X_3)v_k + (A^2 - A^{-2})\rho(X_2)v_k\\ &\qquad - x_3A^{2k+2}\rho(X_2)\rho(X_3)v_k + x_3A^{2k+1}(A^2 - A^{-2})\rho(X_1)v_k.\end{aligned}$$

Using the facts that $\rho(X_3)v_k = \lambda_k v_k$ and $\lambda_k = x_3A^{2k} + x_3^{-1}A^{-2k}$, we obtain

$$\begin{aligned}\rho(X_3)\mathcal{U}_k v_k &= A^{-1}\rho(X_1)\lambda_k v_k + (A^2 - A^{-2})\rho(X_2)v_k\\ &\qquad - x_3A^{2k+2}\rho(X_2)\lambda_k v_k + x_3A^{2k+1}(A^2 - A^{-2})\rho(X_1)v_k\\ &= (x_3A^{2k+3} + x_3^{-1}A^{-2k-1})\rho(X_1)v_k - (x_3^2A^{4k+2} + A^{-2})\rho(X_2)v_k\\ &= \lambda_{k+1}\big(A\rho(X_1) - x_3A^{2k}\rho(X_2)\big)v_k = \lambda_{k+1}\mathcal{U}_k v_k\end{aligned}$$

which shows that $\mathcal{U}_k v_k$ belongs to V_{k+1}.

The proof that $\mathcal{D}_k$ sends V_k to V_{k-1} is similar. $\square$

The indexing of the eigenspaces V_k of $\rho(X_3)$ and of the operators $\mathcal{U}_k$, $\mathcal{D}_k$ depends on our choice of x_3 such that $t_3 = x_3^N + x_3^{-N}$. In particular, replacing x_3 by x_3A^{2l} replaces V_k, $\mathcal{U}_k$, $\mathcal{D}_k$ by V_{k-l}, $\mathcal{U}_{k-l}$ and $\mathcal{D}_{k-l}$, respectively. Similarly, replacing x_3 by x_3^{-1} flips the order and replaces V_k, $\mathcal{U}_k$, $\mathcal{D}_k$ by V_{N-k}, $\mathcal{D}_{N-k}$, $\mathcal{U}_{N-k}$.

Lemma 9. *For every vector $v_k \in V_k$,*

$$\begin{aligned}
\rho(X_1)v_k &= -\frac{x_3^{-1}A^{-2k-1}}{x_3A^{2k} - x_3^{-1}A^{-2k}}\mathcal{U}_k v_k + \frac{x_3A^{2k-1}}{x_3A^{2k} - x_3^{-1}A^{-2k}}\mathcal{D}_k v_k \\
\rho(X_2)v_k &= -\frac{1}{x_3A^{2k} - x_3^{-1}A^{-2k}}\mathcal{U}_k v_k + \frac{1}{x_3A^{2k} - x_3^{-1}A^{-2k}}\mathcal{D}_k v_k \\
\rho(X_3)v_k &= (x_3A^{2k} + x_3^{-1}A^{-2k})v_k.
\end{aligned}$$

Proof. The first two lines are obtained by solving for $\rho(X_1)v_k$ and $\rho(X_3)v_k$ in the definition of $\mathcal{U}_k$ and $\mathcal{D}_k$. We just need to check that the denominators are non-zero. But this immediately follows from the hypothesis that $t_3 = x_3^N + x_3^{-N}$ is different from ±2.

The last equation is just a rephrasing of the definition of the subspace V_k. ☐

Our next computation shows how the operators $\mathcal{U}_k$ and $\mathcal{D}_k$ interact with each other. Note that $\mathcal{U}_k$ sends V_k to V_{k+1}, and that $\mathcal{D}_{k+1}$ sends V_{k+1} back to V_k.

Lemma 10. *For every $v_k \in V_k$,*

$$\mathcal{D}_{k+1}\mathcal{U}_k v_k = -(p + x_3^2A^{4k+2} + x_3^{-2}A^{-4k-2})v_k.$$

Proof. We begin by expanding

$$\begin{aligned}
\mathcal{D}_{k+1}\mathcal{U}_k v_k &= \big(A\rho(X_1) - x_3^{-1}A^{-2k-2}\rho(X_2)\big)\big(A\rho(X_1) - x_3A^{2k}\rho(X_2)\big)v_k \\
&= \big(A^2\rho(X_1^2) - x_3^{-1}A^{-2k-1}\rho(X_2X_1) - x_3A^{2k+1}\rho(X_1X_2) + A^{-2}\rho(X_2^2)\big)v_k.
\end{aligned}$$

The puncture invariant p is defined by the property that $\rho(P) = p\,\mathrm{Id}_V$ for the puncture element

$$P = AX_1X_2X_3 - A^2X_1^2 - A^{-2}X_2^2 - A^2X_3^2 + A^2 + A^{-2}.$$

Therefore, using the fact that $\rho(X_3)v_k = \lambda_k v_k = (x_3A^{2k} + x_3^{-1}A^{-2k})v_k$ and the relation that $AX_1X_2 - A^{-1}X_2X_1 = (A^2 - A^{-2})X_3$,

$$\begin{aligned}
\mathcal{D}_{k+1}\mathcal{U}_k v_k + pv_k &= A^2\rho(X_1^2)v_k - x_3^{-1}A^{-2k-1}\rho(X_2X_1)v_k - x_3A^{2k+1}\rho(X_1X_2)v_k \\
&\qquad + A^{-2}\rho(X_2^2)v_k + A\rho(X_1X_2X_3)v_k - A^2\rho(X_1^2)v_k \\
&\qquad - A^{-2}\rho(X_2^2)v_k - A^2\rho(X_3^2)v_k + (A^2 + A^{-2})v_k \\
&= -(x_3^2A^{4k+2} + x_3^{-2}A^{-4k-2})v_k
\end{aligned}$$

after simplifications. This concludes the proof. ☐

Lemma 11. *Consider the map*

$$\prod_{j=1}^N \mathcal{D}_{k+j} \prod_{j=1}^N \mathcal{U}_{k+N-j} = \mathcal{D}_{k+1}\mathcal{D}_{k+2}\dots\mathcal{D}_{k+N-1}\mathcal{D}_{k+N}\mathcal{U}_{k+N-1}\mathcal{U}_{k+N-2}\dots\mathcal{U}_{k+1}\mathcal{U}_k.$$

For every $v_k \in V_k$,

$$\prod_{j=1}^N \mathcal{D}_{k+j} \prod_{j=1}^N \mathcal{U}_{k+N-j} v_k = (t_1t_2t_3 + t_1^2 + t_2^2)v_k$$

Proof. Note that $\mathcal{U}_{k+j-1}\mathcal{U}_{k+j-2}\dots\mathcal{U}_{k+1}\mathcal{U}_k v_k$ is an element of V_j for every j. By successive applications of Lemma 10, we conclude that

$$\prod_{j=1}^{N}\mathcal{D}_{k+j}\prod_{j=1}^{N}\mathcal{U}_{k+N-j}v_k = -\prod_{j=1}^{N}(p+x_3^2A^{4k+4j+2}+x_3^{-2}A^{-4k-4j-2})v_k.$$

Because A^2 and A^4 are primitive N-roots of unity, the set of powers A^{4k+4j} with $j=1$, 2, …, N is the same as the set of powers A^{2l} with $l=1$, 2, …, N. Therefore

$$\begin{aligned}\prod_{j=1}^{N}(p+x_3^2A^{4k+4j+2}+x_3^{-2}A^{-4k-4j-2}) &= \prod_{l=1}^{N}(p+x_3^2A^{2l}+x_3^{-2}A^{-2l})\\ &= T_N(p)+x_3^{2N}+x_3^{-2N}\\ &= -t_1t_2t_3-t_1^2-t_2^2\end{aligned}$$

by Lemmas 4(b) and 7, remembering that $t_3 = x_3^N + x_3^{-N}$. This completes the proof. □

We now use the fact that the representation $\rho\colon \mathcal{S}_A(S_{1,1}) \to \mathrm{End}(V)$ is irreducible.

Lemma 12. *The space V has dimension N, and all eigenspaces V_k of $\rho(X_3)$ are 1-dimensional.*

More precisely, V admits a basis $\{v_1, v_2, \dots, v_N\}$ where each v_k generates the eigenspace V_k, and where for some $u \neq 0$

$$\mathcal{U}_k v_k = \begin{cases} v_{k+1} & \text{if } 1 \le k \le N-1\\ uv_1 & \text{if } k = N\end{cases}$$

and

$$\mathcal{D}_k v_k = \begin{cases} -\frac{1}{u}(p+x_3^2A^2+x_3^{-2}A^{-2})v_N & \text{if } k=1\\ -(p+x_3^2A^{4k-2}+x_3^{-2}A^{-4k+2})v_{k-1} & \text{if } 2\le k\le N.\end{cases}$$

Proof. The operator $\rho(X_3) \in \mathrm{End}(V)$ admits at least one non-zero eigenvector. Therefore, some V_{k_0} is different from 0.

The map $\prod_{j=1}^{N}\mathcal{U}_{k_0+N-j} = \mathcal{U}_{k_0+N-1}\mathcal{U}_{k_0+N-2}\dots\mathcal{U}_{k_0+1}\mathcal{U}_{k_0}$ of Lemma 11 sends V_{k_0} to V_{k_0}. Because $t_1t_2t_3 + (t_1)^2 + (t_2)^2 \neq 0$ by hypothesis in Theorem 6, Lemma 11 shows that this map is not the 0 map on V_{k_0}. Therefore, there is an eigenvalue $u \neq 0$ and a non-zero eigenvector $v_{k_0} \in V_{k_0}$ such that

$$\mathcal{U}_{k_0+N-1}\mathcal{U}_{k_0+N-2}\dots\mathcal{U}_{k_0+1}\mathcal{U}_{k_0}v_{k_0} = uv_{k_0}.$$

We can now arrange that $k_0 = 1$. Indeed, if we apply $\mathcal{U}_N\mathcal{U}_{N-1}\dots\mathcal{U}_{k_0+1}\mathcal{U}_{k_0}$ to both sides of the above equality and if we set $v_1 = \mathcal{U}_N\mathcal{U}_{N-1}\dots\mathcal{U}_{k_0+1}\mathcal{U}_{k_0}v_{k_0} \in V_{N+1} = V_1$, we obtain that

$$\mathcal{U}_N\mathcal{U}_{N-1}\dots\mathcal{U}_2\mathcal{U}_1v_1 = uv_1.$$

Then, set $v_k = \mathcal{U}_{k-1}\mathcal{U}_{k-2}\dots\mathcal{U}_2\mathcal{U}_1v_1 \in V_k$ for every $k = 1$, 2, …, N. Let W be the N-dimensional linear subspace of V spanned by the vectors v_1, v_2, …, v_N.

By construction,

$$\mathcal{U}_k v_k = \begin{cases} v_{k+1} & \text{if } 1 \le k \le N-1\\ uv_1 & \text{if } k = N.\end{cases}$$

Also, by Lemma 10,

$$\mathcal{D}_k v_k = \begin{cases} -\frac{1}{u}(p + x_3^2 A^2 + x_3^{-2} A^{-2}) v_N & \text{if } k = 1 \\ -(p + x_3^2 A^{4k-2} + x_3^{-2} A^{-4k+2}) v_{k-1} & \text{if } 2 \leq k \leq N. \end{cases}$$

An immediate consequence of these formulas is that, for every k, $\mathcal{U}_k v_k$ and $\mathcal{D}_k v_k$ both belong to W. Lemma 9 then shows that $\rho(X_1)v_k$, $\rho(X_2)v_k$ and $\rho(X_3)v_k$ belong to W. This proves that W is invariant under the image of the representation $\rho\colon \mathcal{S}_A(S_{1,1}) \to \mathrm{End}(V)$. By irreducibility of ρ, it follows that $W = V$. Therefore, V has dimension N, and each eigenspace V_k of $\rho(X_3)$ is the line generated by the vector v_k. □

Combining the above formulas for $\mathcal{U}_k v_k$ and $\mathcal{D}_k v_k$ with Lemma 9 completely determines $\rho(X_1)$ and $\rho(X_2)$ in terms of the parameters p, x_3 and u. Here, p and x_3 are given by the data of Theorem 6. We now determine u in terms of the rest of this data.

Lemma 13.

$$u = -t_1 - x_3^N t_2.$$

Proof. We need to relate the number u to the numbers t_i such that $T_N\big(\rho(X_i)\big) = t_i \,\mathrm{Id}_V$.

By Lemma 9, $\rho(X_1)$ can be decomposed as $\rho(X_1) = \mathcal{U} + \mathcal{D}$ where, in the basis $\{v_1, v_2, \dots, v_N\}$ provided by Lemma 12,

$$\mathcal{U} v_k = -\frac{x_3^{-1} A^{-2k-1}}{x_3 A^{2k} - x_3^{-1} A^{-2k}} \mathcal{U}_k v_k$$
$$\mathcal{D} v_k = \frac{x_3 A^{2k-1}}{x_3 A^{2k} - x_3^{-1} A^{-2k}} \mathcal{D}_k v_k.$$

The crucial property is that $\mathcal{U}$ and $\mathcal{D}$ cyclically permute the eigenspaces V_k in opposite directions, namely $\mathcal{U}$ sends each V_k to V_{k+1} while $\mathcal{D}$ sends V_k to V_{k-1}.

Expanding $T_N\big(\rho(X_1)\big) = T_N(\mathcal{U} + \mathcal{D})$ gives a linear combination of terms of the form $A_1 A_2 \dots A_m$, where each A_i is either $\mathcal{U}$ or $\mathcal{D}$, and where $m \leq N$ is odd. If such a monomial contains n terms that are $\mathcal{U}$ and $m - n$ terms that are $\mathcal{D}$, it sends V_k to V_{k+2n-m}. Therefore, since $T_N\big(\rho(X_1)\big) = t_1 \,\mathrm{Id}_V$, this expansion of $T_N(\mathcal{U} + \mathcal{D})$ contains only terms such that $2n - m = 0 \mod N$. Since m is odd and $0 \leq n \leq m \leq N$, we only have two possibilities: $(m, n) = (N, 0)$ or $(m, n) = (N, N)$. Since the highest degree term of $T_N(x)$ is x^N, this proves that

$$T_N\big(\rho(X_1)\big) = T_N(\mathcal{U} + \mathcal{D}) = \mathcal{U}^N + \mathcal{D}^N.$$

As A^2 is a primitive N-root of unity, $\prod_{k=1}^N (x_3 A^{2k} - x_3^{-1} A^{-2k}) = x_3^N - x_3^{-N}$. It follows that

$$\mathcal{U}^N v_k = \frac{x_3^{-N}}{x_3^N - x_3^{-N}} \mathcal{U}_{k+N-1} \mathcal{U}_{k+N-2} \dots \mathcal{U}_{k+1} \mathcal{U}_k v_k$$
$$= \frac{u x_3^{-N}}{x_3^N - x_3^{-N}} v_k$$

using the expression for $\mathcal{U}_l v_l$ given by Lemma 12.

Similarly,

$$
\begin{aligned}
\mathcal{D}^N v_k &= -\frac{x_3^N}{x_3^N - x_3^{-N}} \mathcal{D}_{k-N+1}\mathcal{D}_{k-N+2}\dots\mathcal{D}_{k-1}\mathcal{D}_k v_k \\
&= +\frac{x_3^N}{x_3^N - x_3^{-N}} \frac{1}{u} \prod_{j=1}^{N} (p + x_3^2 A^{4k-4j+2} + x_3^{-2} A^{-4k+4j-2}) v_k \\
&= -\frac{u^{-1} x_3^N}{x_3^N - x_3^{-N}} (t_1 t_2 t_3 + t_1^2 + t_2^2) v_k
\end{aligned}
$$

where the last equality is proved by the same computation as in the proof of Lemma 11.

Comparing the equalities $T_N\big(\rho(X_1)\big) = t_1 \operatorname{Id}_V$ and $T_N\big(\rho(X_1)\big) = \mathcal{U}^N + \mathcal{D}^N$, it follows that

$$t_1 = -\frac{u x_3^{-N}}{x_3^N - x_3^{-N}} - \frac{u^{-1} x_3^N}{x_3^N - x_3^{-N}} (t_1 t_2 t_3 + t_1^2 + t_2^2).$$

This almost determines u, up to two possibilities. To resolve the ambiguity, we perform similar computations for $T_N\big(\rho(X_2)\big) = t_2 \operatorname{Id}_V$, which give

$$t_2 = \frac{u}{x_3^N - x_3^{-N}} + \frac{u^{-1}}{x_3^N - x_3^{-N}} (t_1 t_2 t_3 + t_1^2 + t_2^2).$$

Combining these expressions for t_1 and t_2 shows that $u = -t_1 - t_2 x_3^N$. □

We now just need to combine Lemmas 9, 12 and 13 to obtain:

Lemma 14.

$$
\rho(X_1) v_k = \begin{cases}
-\frac{x_3^{-1} A^{-3}}{x_3 A^2 - x_3^{-1} A^{-2}} v_2 - \frac{x_3 A (p + x_3^2 A^2 + x_3^{-2} A^{-2})}{(x_3 A^2 - x_3^{-1} A^{-2})(t_1 + t_2 x_3^N)} v_N & \text{if } k = 1 \\
-\frac{x_3^{-1} A^{-2k-1}}{x_3 A^{2k} - x_3^{-1} A^{-2k}} v_{k+1} + \frac{x_3 A^{2k-1} (p + x_3^2 A^{4k-2} + x_3^{-2} A^{-4k+2})}{x_3 A^{2k} - x_3^{-1} A^{-2k}} v_{k-1} & \text{if } 2 \leq k \leq N-1 \\
\frac{x_3^{-1} A^{-1} (t_1 + t_2 x_3^N)}{x_3 - x_3^{-1}} v_1 + \frac{x_3 A^{-1} (p + x_3^2 A^{-2} + x_3^{-2} A^2)}{x_3 - x_3^{-1}} v_{k-1} & \text{if } k = N
\end{cases}
$$

$$
\rho(X_2) v_k = \begin{cases}
-\frac{1}{x_3 A^2 - x_3^{-1} A^{-2}} v_2 - \frac{p + x_3^2 A^2 + x_3^{-2} A^{-2}}{(x_3 A^2 - x_3^{-1} A^{-2})(t_1 + t_2 x_3^N)} v_N & \text{if } k = 1 \\
-\frac{1}{x_3 A^{2k} - x_3^{-1} A^{-2k}} v_{k+1} + \frac{p + x_3^2 A^{4k-2} + x_3^{-2} A^{-4k+2}}{x_3 A^{2k} - x_3^{-1} A^{-2k}} v_{k-1} & \text{if } 2 \leq k \leq N-1 \\
\frac{t_1 + t_2 x_3^N}{x_3 - x_3^{-1}} v_1 + \frac{p + x_3^2 A^{-2} + x_3^{-2} A^{+2}}{x_3 - x_3^{-1}} v_{k-1} & \text{if } k = N
\end{cases}
$$

$$\rho(X_3) v_k = (x_3 A^{2k} + x_3^{-1} A^{-2k}) v_k.$$ □

This proves that, up to isomorphism, the representation ρ is completely determined by the numbers t_1, t_2, x_3 and p. Since x_3 was chosen as an arbitrary number such that $x_3^N + x_3^{-N} = t_3$, it follows that ρ is completely determined by t_1, t_2, t_3, p, namely by its classical shadow $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{1,1})$ and its puncture invariant p.

This concludes the proof of Theorem 6, and therefore of the Uniqueness Theorem 3 in the case of the one-puncture torus $S_{1,1}$. □

4.3. **A more general statement.**

The following statement slightly improves Theorem 6, by providing fewer exceptions.

Theorem 15. *Consider a character $r \in \mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{1,1})$. Setting $t_i = -\operatorname{Tr} r(X_i)$ for the generators X_i of Proposition 5, suppose that at least one of the following conditions fails:*

(1) *$t_i = \pm 2$ for each $i = 1, 2, 3$;*

(2) *$t_i = 0$ for each $i = 1, 2, 3$;*
(3) *one trace t_i is equal to ± 2, the other two t_j are equal to $\pm\frac{2}{\sqrt{3}}\mathrm{i}$, and the signs are such that $t_1t_2t_3 = -\frac{8}{3}$.*

Then, up to isomorphism, there exists a unique representation $\rho\colon \mathcal{S}_A(S_{1,1}) \to \mathrm{End}(V)$ with classical shadow r and puncture invariant p for every $p \in \mathbb{C}$ with

$$T_N(p) = -t_1t_2t_3 - t_1^2 - t_2^2 - t_3^2 + 2.$$

Proof. An easy case-by-case analysis show that, if the hypotheses of Theorem 6 hold, it is possible to cyclically reindex the curves X_1, X_2, X_3 (which does not change their relations) so that $t_3 \neq \pm 2$ and $t_1t_2t_3 + t_1^2 + t_2^2 \neq 0$, namely so that the hypotheses of Theorem 6 are satisfied. Theorem 6 then proves the uniqueness of the representation ρ.

To prove the existence, one can rely on [3, 4] or check by brute force computation that the operators explicitly given in Lemma 14 really satisfy the relations of Proposition 5. □

5. The unpunctured torus $S_{1,0}$

For the unpunctured torus $S_{1,0}$, Bullock and Przytycki show that the skein algebra $\mathcal{S}_A(S_{1,0})$ admits a presentation with the same generators and relations as $\mathcal{S}_A(S_{1,1})$ in Proposition 5, but with the additional relation that $P + A^2 + A^{-2} = 0$. As a consequence, an irreducible representation $\rho\colon \mathcal{S}_A(S_{1,0}) \to \mathrm{End}(V)$ is equivalent to the data of an irreducible representation $\rho\colon \mathcal{S}_A(S_{1,0}) \to \mathrm{End}(V)$ with puncture invariant $p = -A^2 - A^{-2}$. The Generic Uniqueness Theorem 3 then follows from the case of the one-puncture torus, as proved by Theorem 6.

6. The four-puncture sphere $S_{0,4}$

6.1. A presentation for the skein algebra $\mathcal{S}_A(S_{0,4})$.

A presentation of the skein algebra $\mathcal{S}_A(S_{0,4})$ of the four-puncture sphere $S_{0,4}$ can again be found in Bullock-Przytycki [7].



Figure 3. Curves on the four-puncture sphere

Consider the sphere as a "pillowcase" obtained from a rectangle $[0,2] \times [0,1]$ by identifying each point $(0,y)$ to $(1,y)$, each $(x,0)$ to $(2-x,0)$, and each $(x,1)$ to $(2-x,1)$. Identify the four-puncture sphere $S_{0,4}$ to the surface obtained from this pillowcase by removing the four points that are the images of the six points of $[0,2]\times[0,1]$ with integer coordinates.

This enables us to single out several simple closed curves in $S_{0,4}$. The first curve X_1 in the image of the two vertical arcs $\{\frac{1}{2},\frac{3}{2}\}\times[0,1]$. The second curve X_3 is the image of the horizontal line segment $[0,1]\times\{\frac{1}{2}\}$. The third curve is the image of three slope 1 segments, respectively parametrized by $t\mapsto(\frac{1}{2}t,\frac{1}{2}t+\frac{1}{2})$, $t\mapsto(t+\frac{1}{2},t)$ and $t\mapsto(\frac{1}{2}t+\frac{3}{2},\frac{1}{2}t)$ for $0\leq t\leq 1$.

We also consider small loops P_0, P_1, P_2, P_3 going around the punctures, indexed in such a way that for $i=1$, 2, 3, the closed curve X_i separates P_0 and P_i from the other two puncture loops. See Figure 3.

We consider the elements X_1, X_2, X_3, P_0, P_1, P_2, $P_3\in\mathcal{S}_A(S_{0,4})$ represented by these simple closed curves, endowed with the vertical framing.

Proposition 16 ([7]). *The skein algebra $\mathcal{S}_A(S_{0,4})$ of the four-puncture sphere admits a presentation with generators X_1, X_2, X_3, P_0, P_1, P_2, P_3 as above, and with the following relations:*

1. *the P_i are central;*
2. $A^2X_1X_2-A^{-2}X_2X_1=(A^4-A^{-4})X_3+(A^2-A^{-2})(P_0P_3+P_1P_2)$
3. $A^2X_2X_3-A^{-2}X_3X_2=(A^4-A^{-4})X_1+(A^2-A^{-2})(P_0P_1+P_2P_3)$
4. $A^2X_3X_1-A^{-2}X_1X_3=(A^4-A^{-4})X_2+(A^2-A^{-2})(P_0P_2+P_1P_3)$
5. $A^2X_1X_2X_3-A^4X_1^2-A^{-4}X_2^2-A^4X_3^2-A^2(P_0P_1+P_2P_3)X_1-A^{-2}(P_0P_2+P_1P_3)X_2-A^2(P_0P_3+P_1P_2)X_3+(A^2+A^{-2})^2=P_0P_1P_2P_3+P_0^2+P_1^2+P_2^2+P_3^2.$

We will take advantage of the fact that the relations (2–4) are very similar to the relations what we already encountered for the one-puncture torus.

6.2. Reconstructing an irreducible representations of $\mathcal{S}_A(S_{0,4})$ from its invariants.

Let $\rho\colon\mathcal{S}_A(S_{0,4})\to\mathrm{End}(V)$ be an irreducible representation with classical shadow $r\in\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{0,4})$ and puncture invariants p_0, p_1, p_2, p_3. As in the proof of Theorem 6, we want to show that ρ can be reconstructed from these invariants, provided that they are generic enough.

Theorem 17. *Let $\rho\colon\mathcal{S}_A(S_{0,4})\to End(V)$ be an irreducible representation with classical shadow $r\in\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{0,4})$ and puncture invariants p_0, p_1, p_2. $p_3\in\mathbb{C}$. Suppose that,*

$$\operatorname{Tr}r(X_3)\neq\pm2$$

$$\text{and } \operatorname{Tr}r(X_3)\neq T_N(r) \text{ for every solution } r \text{ of the equation}$$

$$\big(r^2+p_0p_3r+p_0^2+p_3^2-4\big)\big(r^2+p_1p_2r+p_1^2+p_2^2-4\big)=0.$$

Then, up to isomorphism, the representation $\rho\colon\mathcal{S}_A(S_{1,1})\to End(V)$ is completely determined by r and p. In addition, V has dimension N.

Theorem 17 proves the Generic Uniqueness Theorem 3 in the case of the four-puncture sphere.

The rest of this section is devoted to the proof of Theorem 17. In particular, we henceforth assume that its hypotheses hold. The proof follows the general lines of the argument used for Theorem 6, but most steps will be more complicated. As a consequence, our reconstruction of the representation ρ from its classical shadow $r\in\mathcal{X}_{\mathrm{SL}_2(\mathbb{C})}(S_{0,4})$ and its puncture invariants p_i will not be as explicit as in that earlier case.

As in the case of the one-puncture torus, set $t_i = -\operatorname{Tr} r(X_i)$, so that $T_N\big(\rho(X_i)\big) = t_i \operatorname{Id}_V$. Also, in view of the relations of Proposition 16, it is convenient to introduce

$$\begin{aligned}
q_1 &= p_0p_1 + p_2p_3\\
q_2 &= p_0p_2 + p_1p_3\\
q_3 &= p_0p_3 + p_1p_2\\
\Delta &= p_0p_1p_2p_3 + p_0^2 + p_1^2 + p_2^2 + p_3^2.
\end{aligned}$$

Choose $x_3 \in \mathbb{C}$, such that $t_3 = x_3^N + x_3^{-N}$. Since $T_N\big(\rho(X_3)\big) = t_3 \operatorname{Id}_V$, all possible eigenvalues of $\rho(X_3)$ are of the form $\lambda_k = x_3A^{4k} + x_3^{-1}A^{-4k}$ for $k = 1, 2, \dots, N-1$. These N numbers λ_k are distinct by our hypothesis that $t_3 \neq \pm 2$. As before, set $V_k = \{v \in V; \rho(X_3)v = \lambda_k v\}$.

The "up" and "down" operators $\mathcal{U}_k$ and $\mathcal{D}_k$ are now given by more complicated formulas.

$$\begin{aligned}
\mathcal{U}_k &= A^2\rho(X_1) - x_3A^{4k}\rho(X_2) + \frac{q_2 + x_3A^{4k+2}q_1}{x_3A^{4k+2} - x_3^{-1}A^{-4k-2}}\operatorname{Id}_V\\
&= A^2\rho(X_1) - x_3A^{4k}\rho(X_2) + \beta_k^+ \operatorname{Id}_V\\
\mathcal{D}_k &= A^2\rho(X_1) - x_3^{-1}A^{-4k}\rho(X_2) + \frac{-q_2 - x_3^{-1}A^{-4k+2}q_1}{x_3A^{4k-2} - x_3^{-1}A^{-4k+2}}\operatorname{Id}_V\\
&= A^2\rho(X_1) - x_3^{-1}A^{-4k}\rho(X_2) + \beta_k^- \operatorname{Id}
\end{aligned}$$

if we set $\beta_k^+ = \frac{q_2+x_3A^{4k+2}q_1}{x_3A^{4k+2}-x_3^{-1}A^{-4k-2}}$ and $\beta_k^- = \frac{-q_2-x_3^{-1}A^{-4k+2}q_1}{x_3A^{4k-2}-x_3^{-1}A^{-4k+2}}$ to simplify further computations.

Lemma 18. *For every $v_k \in V_k$,*

$$\begin{aligned}
\rho(X_1)v_k &= -\frac{x_3^{-1}A^{-4k-2}}{x_3A^{4k} - x_3^{-1}A^{-4k}}\mathcal{U}_kv_k + \frac{x_3A^{4k-2}}{x_3A^{4k} - x_3^{-1}A^{-4k}}\mathcal{D}_kv_k\\
&\qquad + \frac{x_3^{-1}A^{-4k-2}\beta_k^+ - x_3A^{4k-2}\beta_k^-}{x_3A^{4k} - x_3^{-1}A^{-4k}}v_k\\
\rho(X_2)v_k &= -\frac{1}{x_3A^{4k} - x_3^{-1}A^{-4k}}\mathcal{U}_kv_k + \frac{1}{x_3A^{4k} - x_3^{-1}A^{-4k}}\mathcal{D}_kv_k\\
&\qquad + \frac{\beta_k^+ - \beta_k^-}{x_3A^{4k} - x_3^{-1}A^{-4k}}v_k\\
\rho(X_3)v_k &= (x_3A^{4k} + x_3^{-1}A^{-4k})v_k.
\end{aligned}$$

□

Lemma 19. *The operator $\mathcal{U}_k$ sends each subspace V_k to V_{k+1}, and $\mathcal{D}_k$ sends V_k to V_{k-1}.*

Proof. Given a vector $v_k \in V_k$, we want to show that $\rho(X_3)\mathcal{U}_kv_k = \lambda_{k+1}\mathcal{U}_kv_k$. As in the proof of Lemma 8, we expand

$$\begin{aligned}
\rho(X_3)\mathcal{U}_kv_k &= A^2\rho(X_3X_1)v_k - x_3A^{4k}\rho(X_3X_2)v_k + \beta_k^+\rho(X_3)v_k\\
&= A^{-2}\rho(X_1)\rho(X_3)v_k + (A^4 - A^{-4})\rho(X_2)v_k + (A^2 - A^{-2})q_2v_k\\
&\qquad - x_3A^{4k+4}\rho(X_2)\rho(X_3)v_k + x_3A^{4k+2}(A^4 - A^{-4})\rho(X_1)v_k\\
&\qquad + x_3A^{4k+2}(A^2 - A^{-2})q_1v_k + \beta_k^+\rho(X_3)v_k
\end{aligned}$$

using the relations of Proposition 16.

Since $\rho(X_3)v_k = \lambda_k v_k$ and $\lambda_k = (x_3A^{4k} + x_3^{-1}A^{-4k})v_k$, it follows that

$$\begin{aligned}\rho(X_3)\mathcal{U}_kv_k &= A^{-2}\rho(X_1)\lambda_kv_k + (A^4 - A^{-4})\rho(X_2)v_k + (A^2 - A^{-2})q_2v_k\\ &\qquad - x_3A^{4k+4}\rho(X_2)\lambda_kv_k + x_3A^{4k+2}(A^4 - A^{-4})\rho(X_1)v_k\\ &\qquad + x_3A^{4k+2}(A^2 - A^{-2})q_1v_k + \beta_k^+\lambda_kv_k\\ &= (x_3A^{4k+6} + x_3^{-1}A^{-4k-2})\rho(X_1)v_k - (x_3^2A^{8k+4} + A^{-4})\rho(X_2)v_k\\ &\qquad + \beta_k^+(x_3A^{4k+4} + x_3^{-1}A^{-4k-4})v_k\\ &= \lambda_{k+1}\big(A^2\rho(X_1) - x_3A^{4k}\rho(X_2) + \beta_k^+\big)v_k = \lambda_{k+1}\mathcal{U}_kv_k\end{aligned}$$

which shows that $\mathcal{U}_kv_k$ belongs to V_{k+1}.

The proof that $\mathcal{D}_k$ sends V_k to V_{k-1} is very similar. □

We now have the equivalent of Lemma 10.

Lemma 20. *For every $v_k \in V_k$,*

$$\mathcal{D}_{k+1}\mathcal{U}_kv_k = R_kv_k$$

where

$$R_k = -(\Delta - 2 + x_3^2A^{8k+4} + x_3^{-2}A^{-8k-4} + (x_3A^{4k+2} + x_3^{-1}A^{-4k-2})q_3 - \beta_{k+1}^-\beta_k^+)$$

Proof. We begin by expanding

$$\begin{aligned}\mathcal{D}_{k+1}\mathcal{U}_kv_k &= (A^2\rho(X_1) - x_3^{-1}A^{-4k-4}\rho(X_2) + \beta_{k+1}^-\,\mathrm{Id}_V)(A^2\rho(X_1) - x_3A^{4k}\rho(X_2) + \beta_k^+\,\mathrm{Id}_V)v_k\\ &= A^4\rho(X_1^2)v_k - x_3^{-1}A^{-4k-2}\rho(X_2X_1)v_k + A^2\beta_{k+1}^-\rho(X_1)v_k\\ &\qquad - x_3A^{4k+2}\rho(X_1X_2)v_k + A^{-4}\rho(X_2^2)v_k - x_3A^{4k}\beta_{k+1}^-\rho(X_2)v + k\\ &\qquad + A^2\beta_k^+\rho(X_1)v_k - x_3^{-1}A^{-4k-4}\beta_k^+\rho(X_2)v_k + \beta_{k+1}^-\beta_k^+v_k\end{aligned}$$

Using Relation (5) of Proposition 16,

$$\begin{aligned}\mathcal{D}_{k+1}\mathcal{U}_kv_k + \Delta v_k &= A^4\rho(X_1^2)v_k - x_3^{-1}A^{-4k-2}\rho(X_2X_1)v_k + A^2\beta_{k+1}^-\rho(X_1)v_k\\ &\qquad - x_3A^{4k+2}\rho(X_1X_2)v_k + A^{-4}\rho(X_2^2)v_k - x_3A^{4k}\beta_{k+1}^-\rho(X_2)v_k\\ &\qquad + A^2\beta_k^+\rho(X_1)v_k - x_3^{-1}A^{-4k-4}\beta_k^+\rho(X_2)v_k + \beta_{k+1}^-\beta_k^+v_k\\ &\qquad + A^2\rho(X_1X_2X_3)v_k - A^4\rho(X_1^2)v_k - A^{-4}\rho(X_2^2) - A^4\rho(X_3^2)v_k\\ &\qquad - A^2q_1\rho(X_1)v_k - A^{-2}q_2\rho(X_2) - A^2q_3\rho(X_3)v_k + (A^2 + A^{-2})^2v_k\\ &= -x_3^{-1}A^{-4k-2}\rho(X_2X_1)v_k + A^2\beta_{k+1}^-\rho(X_1)v_k - x_3A^{4k+2}\rho(X_1X_2)v_k\\ &\qquad - x_3A^{4k}\beta_{k+1}^-\rho(X_2)v_k + A^2\beta_k^+\rho(X_1)v_k - x_3^{-1}A^{-4k-4}\beta_k^+\rho(X_2)v_k\\ &\qquad + \beta_{k+1}^-\beta_k^+v_k + A^2\rho(X_1X_2X_3)v_k - A^4\rho(X_3^2) - A^2q_1\rho(X_1)v_k\\ &\qquad - A^{-2}q_2\rho(X_2)v_k - A^2q_3\rho(X_3)v_k + (A^2 + A^{-2})^2v_k.\end{aligned}$$

We simplify this complicated expression in a few steps. We first observe that all terms with $\rho(X_1)$ and $\rho(X_2)$ cancel each other out. Indeed,

$$A^2\beta_{k+1}^-\rho(X_1)v_k + A^2\beta_k^+\rho(X_1)v_k - A^2q_1\rho(X_1)v_k = A^2(\beta_{k+1}^- + \beta_k^+ - q_1)\rho(X_1)v_k = 0$$

and

$$\begin{aligned}-x_3A^{4k}\beta^-_{k+1}\rho(X_2)v_k - x_3^{-1}A^{-4k-4}\beta^+_k\rho(X_2)v_k - A^{-2}q_2\rho(X_2)v_k\\ = -A^{-2}(x_3A^{4k+2}\beta^-_{k+1} + x_3^{-1}A^{-4k-2}\beta^+_k + q_2)\rho(X_2)v_k = 0,\end{aligned}$$

so that we are left with

$$\begin{aligned}\mathcal{D}_{k+1}\mathcal{U}_kv_k + \Delta v_k = -x_3^{-1}A^{-4k-2}\rho(X_2X_1)v_k - x_3A^{4k+2}\rho(X_1X_2)v_k + \beta^-_{k+1}\beta^+_kv_k\\ + A^2\rho(X_1X_2X_3)v_k - A^4\rho(X_3^2)v_k - A^2q_3\rho(X_3)v_k + (A^2+A^{-2})^2v_k.\end{aligned}$$

We now use the fact that v_k is an eigenvalue of $\rho(X_3)$:

$$A^2\rho(X_1X_2X_3)v_k = A^2\rho(X_1X_2)\lambda_kv_k = A^2(x_3A^{4k} + x_3^{-1}A^{-4k})\rho(X_1X_2)v_k.$$

Therefore

$$\begin{aligned}\mathcal{D}_{k+1}\mathcal{U}_kv_k + \Delta v_k &= -x_3^{-1}A^{-4k-2}\rho(X_2X_1)v_k + x_3^{-1}A^{-4k+2}\rho(X_1X_2)v_k + \beta^-_{k+1}\beta^+_kv_k\\ &\quad - A^4\rho(X_3^2)v_k - A^2q_3\rho(X_3)v_k + (A^2+A^{-2})^2v_k\\ &= x_3^{-1}A^{-4k}(A^4 - A^{-4})\rho(X_3)v_k + (A^2-A^{-2})q_3v_k + \beta^-_{k+1}\beta^+_kv_k\\ &\quad - A^4\rho(X_3^2)v_k - A^2q_3\rho(X_3)v_k + (A^2+A^{-2})^2v_k\end{aligned}$$

after using Relation (1) of Proposition 16. The formula of Lemma 20 then follows from a last application of the property that $\rho(X_3)v_k = (x_3A^{4k} + x_3^{-1}A^{-4k})v_k$. □

We now turn to the analogue of Lemma 11, and consider the product

$$\prod_{j=1}^N \mathcal{D}_{k+j}\prod_{j=1}^N \mathcal{U}_{k+N-j} = \mathcal{D}_{k+1}\mathcal{D}_{k+2}\dots\mathcal{D}_{k+N-1}\mathcal{D}_{k+N}\mathcal{U}_{k+N-1}\mathcal{U}_{k+N-2}\dots\mathcal{U}_{k+1}\mathcal{U}_k.$$

Since the output of Lemma 20 is more complicated than that of Lemma 10, the computation will be more elaborate. We begin with a simple step.

Lemma 21. *For every $v_k \in V_k$,*

$$\prod_{j=1}^N \mathcal{D}_{k+j}\prod_{j=1}^N \mathcal{U}_{k+l-j}\, v_k = \prod_{j=1}^N R_j\, v_k$$

where R_j is as in Lemma 20.

Proof. As in the proof of Lemma 11, this follows from Lemma 20. □

We next tackle the product of Lemma 21.

Lemma 22.

$$\prod_{j=1}^N R_j = -\frac{\big(t_3 - T_N(r_0)\big)\big(t_3 - T_N(r_1)\big)\big(t_3 - T_N(r_2)\big)\big(t_3 - T_N(r_3)\big)}{t_3^2 - 4}$$

where r_0, r_1, r_2 and r_3 are the solutions of the equation

$$\big(r^2 + p_0p_3r + p_0^2 + p_3^2 - 4\big)\big(r^2 + p_1p_2r + p_1^2 + p_2^2 - 4\big) = 0.$$

Proof. After substituting back the values of $\beta_j^+ = \frac{q_2 + x_3 A^{4j+2} q_1}{x_3 A^{4j+2} - x_3^{-1} A^{-4j-2}}$, $\beta_j^- = \frac{-q_2 - x_3^{-1} A^{-4j+2} q_1}{x_3 A^{4j-2} - x_3^{-1} A^{-4j+2}}$ and $\Delta = p_0 p_1 p_2 p_3 + p_0^2 + p_1^2 + p_2^2 + p_3^2$ and expanding,

$$R_j = -(\Delta - 2 + x_3^2 A^{8j+4} + x_3^{-2} A^{-8j-4} + (x_3 A^{4j+2} + x_3^{-1} A^{-4j-2}) q_3 - \beta_{j+1}^- \beta_j^+)$$

can be factored as $R_j = -S_j S_j' / S_j''$ where

$$\begin{aligned} S_j &= \left(x_3^{-2} A^{-8j-4} + p_0 p_3 x_3^{-1} A^{-4j-2} + \left(p_0^2 + p_3^2 - 2\right) + p_0 p_3 x_3 A^{4j+2} + x_3^2 A^{8j+4}\right) \\ S_j' &= \left(x_3^{-2} A^{-8j-4} + p_1 p_2 x_3^{-1} A^{-4j-2} + \left(p_1^2 + p_2^2 - 2\right) + p_1 p_2 x_3 A^{4j+2} + x_3^2 A^{8j+4}\right) \\ \text{and } S_j'' &= \left(x_3 A^{4j+2} - x_3^{-1} A^{-4j-2}\right)^2 . \end{aligned}$$

The term S_j looks nicer in terms of $u = x_3 A^{4j+2}$. Then,

$$\begin{aligned} S_j &= u^{-2} + u^2 + p_0 p_3 u + p_0 p_3 u^{-1} + p_0^2 + p_3^2 - 2 \\ &= (u + u^{-1})^2 + p_0 p_3 (u + u^{-1}) + p_0^2 + p_3^2 - 4 \\ &= (u + u^{-1} - r_0)(u + u^{-1} - r_3) \\ &= (r_0 - x_3 A^{4j+2} - x_3^{-1} A^{-4j-2})(r_3 - x_3 A^{4j+2} - x_3^{-1} A^{-4j-2}) \end{aligned}$$

where r_0 and r_3 are the solutions of the equation $r^2 + p_0 p_3 r + p_0^2 + p_3^2 - 4 = 0$. Note that r_0 and r_3 do not depend on j. Lemma 4(b) then shows that

$$\prod_{j-1}^{N} S_j = \left(T_N(r_0) - x_3^N - x_3^{-N}\right)\left(T_N(r_3) - x_3^N - x_3^{-N}\right) = \left(T_N(r_0) - t_3\right)\left(T_N(r_3) - t_3\right).$$

Similarly,

$$\prod_{j-1}^{N} S_j' = \left(T_N(r_1) - t_3\right)\left(T_N(r_2) - t_3\right).$$

where r_1 and r_2 are the solutions of the equation $r^2 + p_1 p_2 r + p_1^2 + p_2^2 - 4 = 0$.

Finally

$$\prod_{j-1}^{N} S_j'' = \prod_{j-1}^{N} \left(x_3 A^{4j+2} - x_3^{-1} A^{-4j-2}\right)^2 = \left(x_3^N - x_3^{-N}\right)^2 = t_3^2 - 4,$$

which concludes the computation. □

Lemma 23. *The space V has dimension N, and all eigenspaces V_k of $\rho(X_3)$ are 1-dimensional.*

More precisely, V admits a basis $\{v_1, v_2, \ldots, v_N\}$ where each v_k generates the eigenspace V_k, and where for some $u \neq 0$

$$\mathcal{U}_k v_k = \begin{cases} v_{k+1} & \text{if } 1 \le k \le N-1 \\ u v_1 & \text{if } k = N \end{cases}$$

and

$$\mathcal{D}_k v_k = \begin{cases} \frac{1}{u} R_N v_N & \text{if } k = 1 \\ R_{k-1} v_{k-1} & \text{if } 2 \le k \le N. \end{cases}$$

where R_k is defined as in Lemma 20.

Proof. By the combination of Lemmas 21 and 22, and by hypothesis on t_3, the product $\prod_{j=1}^{N} \mathcal{D}_{k+j} \prod_{j=1}^{N} \mathcal{U}_{k+l-j}$ is different from 0. The proof is then identical to that of Lemma 12. □

Lemma 24. *The number u occurring in Lemma 23 is completely determined by x_3, t_1, t_2 and the puncture invariants p_i.*

Proof. Although we cannot be as specific as in the proof of Lemma 12, we will follow a very similar argument.

By Lemma 18, we can write $\rho(X_1)$ as a sum $\rho(X_1) = \mathcal{U} + \mathcal{D} + \mathcal{I}$ where, in the basis $\{v_1, v_2, \dots, v_N\}$ of Lemma 23,

$$\begin{aligned}
\mathcal{U}v_k &= -\frac{x_3^{-1}A^{-4k-2}}{x_3A^{4k} - x_3^{-1}A^{-4k}}\mathcal{U}_k v_k \\
\mathcal{D}v_k &= \frac{x_3A^{4k-2}}{x_3A^{4k} - x_3^{-1}A^{-4k}}\mathcal{D}_k v_k \\
\mathcal{I}v_k &= \frac{x_3^{-1}A^{-4k-2}\beta_k^+ - x_3A^{4k-2}\beta_k^-}{x_3A^{4k} - x_3^{-1}A^{-4k}}v_k.
\end{aligned}$$

The key observation is that, by the combination of Lemma 23 with the above definition, $\mathcal{U}v_k = a_k v_{k+1}$ where a_k depends only on x_3 if $1 \leq k \leq N-1$, and where a_N is u times a quantity depending only on x_3. Similarly, $\mathcal{D}v_k = b_k v_{k-1}$ where b_k depends only on x_3 and the puncture invariants p_i if $2 \leq k \leq N$, and where b_1 is $\frac{1}{u}$ times a quantity depending only on x_3 and the p_i. And $\mathcal{I}_k = c_k v_k$ where c_k depends only on x_3 and the p_i.

If we expand $T_N\big(\rho(X_1)\big) = T_N(\mathcal{U} + \mathcal{D} + \mathcal{I})$, we obtain a linear combination of terms $A_1A_2\dots A_m$ with $m \leq N$, where each A_i is equal to $\mathcal{U}$, $\mathcal{D}$ or $\mathcal{I}$. Since $T_N\big(\rho(X_1)\big) = t_1 \operatorname{Id}_V$ the only monomials that do not cancel out in this linear combination are those for which the line $A_1A_2\dots A_{m-1}A_mV_1$ is equal to V_1.

For most such monomials $A_1A_2\dots A_m$ with a nontrivial contribution to $T_N\big(\rho(X_1)\big)$, the sequence of lines V_1, A_mV_1, $A_{m-1}A_mV_1$, $\dots$, $A_2\dots A_{m-1}A_mV_1$, $A_1A_2\dots A_{m-1}A_mV_1 = V_1$ switches as many times from V_1 to V_N as it does from V_N to V_1. This implies that, when we compute $A_1A_2\dots A_mv_1$, any term u is balanced by a term $\frac{1}{u}$ and conversely, so that $A_1A_2\dots A_mv_1$ is equal to v_1 times a scalar depending only on x_3 and the p_i.

Because $m \leq N$, there are exactly two exceptions to this property, namely $A_1A_2\dots A_m = \mathcal{U}^N$ and $\mathcal{D}^N$. These two exceptions occur with coefficient 1 in the expression of $T_N\big(\rho(X_1)\big)$ since the highest degree of $T_N(x)$ is x^N. Also,

$$\begin{aligned}
\mathcal{U}^N v_1 &= -u\prod_{k=1}^N \frac{x_3^{-1}A^{-4k-2}}{x_3A^{4k} - x_3^{-1}A^{-4k}}\,v_1 = -u\frac{x_3^{-N}}{x_3^N - x_3^{-N}}\,v_1 \\
\mathcal{D}^N v_1 &= u^{-1}\prod_{k=1}^N \frac{x_3A^{4k-2}}{x_3A^{4k} - x_3^{-1}A^{-4k}}R_{k-1}\,v_1 = u^{-1}\frac{x_3^N}{x_3^N - x_3^{-N}}\prod_{k=1}^N R_k\,v_1
\end{aligned}$$

by Lemma 23.

Since $T_N\big(\rho(X_1)\big) = t_1 \operatorname{Id}_V$, the conclusion of this discussion is that

$$t_1 = -u\frac{x_3^{-N}}{x_3^N - x_3^{-N}} + u^{-1}\frac{x_3^N}{x_3^N - x_3^{-N}}\prod_{k=1}^N R_k + f(x_3, p_0, p_1, p_2, p_3)$$

for an explicit function $f(x_3, p_0, p_1, p_2, p_3)$ of x_3 and of the puncture invariants p_i.

The same argument applied to $T_N\big(\rho(X_2)\big) = t_2 \,\mathrm{Id}_V$ gives that

$$t_2 = -u\frac{1}{x_3^N - x_3^{-N}} + u^{-1}\frac{1}{x_3^N - x_3^{-N}}\prod_{k=1}^N R_k + g(x_3, p_0, p_1, p_2, p_3)$$

for another function $g(x_3, p_0, p_1, p_2, p_3)$ of x_3 and of the puncture invariants p_i.

Then

$$u = t_1 - x_3^N t_2 - f(x_3, p_0, p_1, p_2, p_3) + x_3^N g(x_3, p_0, p_1, p_2, p_3). \qquad \square$$

The combination of Lemmas 18, 23 and 24 then shows that, after isomorphism of ρ, the operators $\rho(X_1)$, $\rho(X_2)$ and $\rho(X_3)$ are completely determined by the t_i and p_j. We are not able to give expressions as explicit as in Lemma 14, but this is enough to prove Theorem 17. ☐

REFERENCES

[1] John W. Barrett. Skein spaces and spin structures. *Math. Proc. Cambridge Philos. Soc.*, 126(2):267–275, 1999.

[2] Francis Bonahon and Helen Wong. Representations of the Kauffman skein algebra I: invariants and miraculous cancellations. preprint, `arXiv:1206.1638`, 2012.

[3] Francis Bonahon and Helen Wong. Representations of the Kauffman skein algebra II: punctured surfaces. preprint, `arXiv:1206.1639`, 2012.

[4] Francis Bonahon and Helen Wong. Representations of the Kauffman skein algebra III: closed surfaces and canonicity. preprint, 2015.

[5] Doug Bullock. Estimating a skein module with $\mathrm{SL}_2(\mathbf{C})$ characters. *Proc. Amer. Math. Soc.*, 125(6):1835–1839, 1997.

[6] Doug Bullock, Charles Frohman, and Joanna Kania-Bartoszyńska. The Kauffman bracket skein as an algebra of observables. *Proc. Amer. Math. Soc.*, 130(8):2479–2485 (electronic), 2002.

[7] Doug Bullock and Józef H. Przytycki. Multiplicative structure of Kauffman bracket skein module quantizations. *Proc. Amer. Math. Soc.*, 128(3):923–931, 2000.

[8] Miroslav Havlíček and Severin Pošta. On the classification of irreducible finite-dimensional representations of $U_q'(\mathrm{so}_3)$ algebra. *J. Math. Phys.*, 42(1):472–500, 2001.

[9] Józef H. Przytycki and Adam S. Sikora. On skein algebras and $\mathrm{Sl}_2(\mathbf{C})$-character varieties. *Topology*, 39(1):115–148, 2000.

[10] Vladimir G. Turaev. Skein quantization of Poisson algebras of loops on surfaces. *Ann. Sci. École Norm. Sup. (4)*, 24(6):635–704, 1991.

E-mail address: `greanvert@gmail.com`